The Noble Element Simulation Technique v2

Jacob Cutter, University of California, Davis
For the NEST Collaboration
NorCal HEP-EX, December 2, 2017

Noble Element TPCs



- Noble elements such as **xenon (Xe)** serve as high quality detection media for reconstructing particle interactions.
- **Dual-phase time projection chambers (TPCs)** are a common example, where energy reconstruction is done using both scintillation and ionization channels.
- It is important to model light and charge yields for a variety of interaction types.

Light and Charge Production in Xenon

- The actual microphysics of scintillation and ionization pathways is complicated, and for Xe we cannot completely track the recoil cascades from first principles.

- NEST (**N**oble **E**lement **S**imulation **T**echnique) began as a semi-empirical model that traced these pathways.

- Precision data has constrained the model to purely empirical fits of yield data.

Lenardo et al.



Motivations for NEST

- Having simulations of yields as a function of electric field is crucial for optimizing detector designs and operational parameters.
- NEST guides not only the process of detector planning, but also informs the data analysis during runs and provides crucial cross-checks.
- Having well-understood background and signal models is crucial for low background experiments and rare event searches.

10 GeV/c^2 WIMP 1000 GeV/c^2 WIMP



LZ TDR (Mount et al.)

Enter NEST v2.0

- The NEST v2.0 is the latest standalone package, which can be used both as a library and as a command-line tool for quick calculations.
- NEST v2 compiles without dependencies on GEANT4 or ROOT, making it a faster and more accessible tool that works right out of the box.
- There are models for various interaction types (each with a succinct formula):
  - Compton scatters and beta decays
  - Photo-absorption
  - ^{83m}Kr

  } Electron recoils (ER)

  - Xe, other heavy nuclear recoils (e.g. ^{206}Pb)
  - Neutrons
  - Alphas

  } Nuclear recoils (NR)

Validation Campaign

Light

Charge

ER







NR







Conclusions

- NEST v2 will be available as a public beta version by the end of this year.
- Future goals include the addition of recombination fluctuations to the physics models, an optional GEANT4 integration for full detector simulations, and a web tool for quick calculations.
- We plan to use the new empirical models as a point of comparison for an eventual first-principles atomic physics model, NEST v3.

NEST Collaboration

UC DAVIS
UNIVERSITY OF CALIFORNIA

Mani Tripathi
Aaron Manalaysay
Jacob Cutter

UNIVERSITY AT ALBANY
State University of New York

Matthew Szydagis
Jack Genovesi
Madison Wyman
Emily Magnus

UNIVERSITY OF MARYLAND
18 56

Jon Balajthy
Carter Hall

UC San Diego

Kaixuan Ni
Junying Huang

MEPhI
NATIONAL RESEARCH NUCLEAR UNIVERSITY • MOSCOW ENGINEERING PHYSICS INSTITUTE • MEPhI •

Ekaterina Kozlova

Lawrence Livermore National Laboratory

Jason Brodsky
Kareem Kazkaz
Brian Lenardo

Berkeley
UNIVERSITY OF CALIFORNIA

Dan McKinsey
Vetri Velan

Thank you!

Questions?